


WHAT IS CLAIMED IS:

1. A fragrance delivery system comprising:
- A) from 0.01% by weight, of a pro-fragrance component which comprises pro-fragrances or pro-accords selected from at least two of the following:
- aldehyde and ketone releasing pro-fragrances, preferably an oxazolidine pro-fragrances;
 - β -amino pro-fragrances; and
 - orthoester pro-accords; and
- B) the balance carriers and other adjunct ingredients.
2. A composition according to Claim 1 further comprising from 1% by weight, a fragrance raw material component comprising:
- optionally at least 1% by weight, of a mixture of one or more base note fragrances;
 - optionally at least 1% by weight, of a mixture of one or more top or middle note fragrances;
 - optionally the balance carriers, fixatives, and other adjunct ingredients.
3. A fragrance raw material delivery system comprising:
- A) from 0.01% by weight, of a pro-fragrance component comprising:
- optionally at least 0.01% by weight, of an aldehyde or ketone releasing pro-fragrance component, said pro-fragrance having the formula:



wherein said pro-fragrance or pro-acord releases an aldehyde or a ketone fragrance raw material, wherein X is oxygen or sulfur; R is:

- $\text{C}_6\text{-C}_{22}$ substituted or unsubstituted linear alkyl;
- $\text{C}_6\text{-C}_{22}$ substituted or unsubstituted branched alkyl;
- $\text{C}_6\text{-C}_{22}$ substituted or unsubstituted linear alkenyl;
- $\text{C}_6\text{-C}_{22}$ substituted or unsubstituted branched alkenyl;

FOI 2007-0467660

Nothing has
to be included;
all components are
optional


- v) C₆-C₂₂ substituted or unsubstituted cycloalkyl;
 vi) C₆-C₂₂ substituted or unsubstituted branched cycloalkyl;
 vii) C₆-C₂₂ substituted or unsubstituted cycloalkenyl;
 viii) C₆-C₂₂ substituted or unsubstituted branched cycloalkenyl;
 ix) C₆-C₂₂ substituted or unsubstituted aryl;
 x) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
 xii) and mixtures thereof;

R¹ is:

- i) hydrogen;
 ii) C₁-C₁₀ substituted or unsubstituted linear alkyl;
 iii) C₃-C₁₀ substituted or unsubstituted branched alkyl;
 iv) C₂-C₁₀ substituted or unsubstituted linear alkenyl;
 v) C₃-C₁₀ substituted or unsubstituted branched alkenyl;
 vi) C₃-C₁₅ substituted or unsubstituted cycloalkyl;
 vii) C₄-C₁₅ substituted or unsubstituted branched cycloalkyl;
 viii) C₄-C₁₅ substituted or unsubstituted cycloalkenyl;
 ix) C₅-C₁₅ substituted or unsubstituted branched cycloalkenyl;
 x) C₆-C₁₅ substituted or unsubstituted aryl;
 xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 xii) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;

R and R¹ can be taken together to form a substituted or unsubstituted ring having in the ring from 3 to 10 carbon atoms; and each R², R³, R⁶ and each R⁴ and R⁵ pair are independently:

- i) R¹;
 ii) hydroxyl;
 iii) a carbonyl comprising unit having the formula:



wherein R⁷ is:

- a) -OH;
 b) -OR⁸ wherein R⁸ is hydrogen, C₁-C₁₅ substituted or unsubstituted linear alkyl, C₁-C₁₅ substituted or unsubstituted branched alkyl, C₂-C₂₂ substituted or unsubstituted linear alkenyl, C₃-C₂₂ substituted or

unsubstituted branched alkenyl, or mixtures thereof; or
 R^8 is M, wherein M is a water soluble cation of
 sufficient charge to render neutrality;

- c) $-N(R^9)_2$ wherein R^9 is hydrogen, C_1 - C_6 substituted or
 unsubstituted linear alkyl, C_3 - C_6 substituted or
 unsubstituted branched alkyl, or mixtures thereof;
- d) C_1 - C_{22} substituted or unsubstituted linear alkyl;
- e) C_1 - C_{22} substituted or unsubstituted branched alkyl;
- f) C_2 - C_{22} substituted or unsubstituted linear alkenyl;
- g) C_3 - C_{22} substituted or unsubstituted branched alkenyl;
- h) C_3 - C_{22} substituted or unsubstituted cycloalkyl;
- i) C_6 - C_{22} substituted or unsubstituted aryl;
- j) C_6 - C_{22} substituted or unsubstituted heterocyclicalkyl;
- k) C_6 - C_{22} substituted or unsubstituted heterocyclicalkenyl;

the index x is from 0 to 22;

- iv) alkyleneoxy units having the formula:



wherein each R^{10} , R^{11} , and R^{12} is independently;

- a) hydrogen;
- b) $-OH$;
- c) C_1 - C_4 alkyl;
- d) or mixtures thereof;

R^{13} is:

- a) hydrogen;
- b) C_1 - C_4 alkyl;
- c) or mixtures thereof;

R^{14} is:

- a) hydrogen;
- b) C_1 - C_4 alkyl;
- c) or mixtures thereof;

R^{10} and R^{11} can be taken together to form a C_3 - C_6 spiroannulated
 ring, carbonyl unit, or mixtures thereof; y has the value from 0 to
 10, z has the value from 1 to 50;


v) any two R^2 , R^3 , R^4 , R^5 , or R^6 units can be taken together to form:

- a) a carbonyl moiety;
- b) a C_3 - C_6 spiroannulated ring;
- c) a heterocyclic aromatic ring comprising from 5 to 7 atoms;
- d) a non-heterocyclic aromatic ring comprising from 5 to 7 atoms;
- e) a heterocyclic ring comprising from 5 to 7 atoms;
- f) a non-heterocyclic ring comprising from 5 to 7 atoms;
- g) or mixtures thereof;


vi) and mixtures thereof; and

the index n is an integer from 1 to 3;

- b) optionally at least 0.01% by weight, of an β -amino pro-fragrance component, said pro-fragrance having the formula:




wherein G^1 is C_1 - C_4 alkyl, -CN, -C(O) Y^1 , -CO $_2Y^1$, Y^2 , and mixtures thereof; G^2 is C_1 - C_4 alkyl, -CN, -C(O) Y^1 , -CO $_2Y^1$, Y^2 , and mixtures thereof; Y^1 and Y^2 are each independently C_1 - C_4 alkyl, or a unit having the formula:



wherein R^6 , R^7 , and R^8 are each independently hydrogen, C_1 - C_4 alkyl, and mixtures thereof; R^6 and R^7 can be taken together to form an exocyclic double bond with the ring; any two R^6 and R^7 , or an R^6 and R^7 with an R^8 can be taken together to form an endocyclic double bond within the ring; two or more R^6 , R^7 , and R^8 units may be taken together


to form one or more C₃-C₇ fused rings, bicyclic rings, or spiroannular rings; m is from 1 to 5;

provided one G¹ or G² is -C(O)Y¹, -CO₂Y¹, or -CN; R and R¹ are each independently hydrogen, C₁-C₂₂ substituted or unsubstituted, branched or unbranched alkyl, C₂-C₂₂ substituted or unsubstituted, branched or unbranched alkenyl, C₂-C₂₀ substituted or unsubstituted, branched or unbranched hydroxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₃-C₂₀ substituted or unsubstituted cycloalkyl, alkyleneoxy units having the formula:




wherein R⁴ is C₂-C₄ alkylene, R⁵ is hydrogen, C₁-C₄ alkyl, and mixtures thereof, x is from 1 to 6; C₆-C₂₀ aryl, C₃-C₂₀ heteroaryl comprising one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof; R and R¹ can be taken together to form one or more aromatic or non-aromatic, heterocyclic or non-heterocyclic, single rings, fused rings, bicyclo rings, spiroannulated rings, or mixtures thereof, said rings comprising from 3 to 20 carbon atoms and one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof;

c) optionally at least 0.01% by weight, of an orthoester pro-accord having the formula:



wherein R is hydrogen, C₁-C₈ linear alkyl, C₄-C₂₀ branched alkyl, C₆-C₂₀ cyclic alkyl, C₆-C₂₀ branched cyclic alkyl, C₆-C₂₀ linear alkenyl, C₆-C₂₀ branched alkenyl, C₆-C₂₀ cyclic alkenyl, C₆-C₂₀ branched cyclic alkenyl, C₆-C₂₀ substituted or unsubstituted aryl, and mixtures thereof; R¹, R² and R³ are independently C₁-C₂₀ linear, branched, or substituted alkyl; C₂-C₂₀ linear, branched, or substituted alkenyl; C₅-C₂₀ substituted or unsubstituted cyclic alkyl; C₆-C₂₀ substituted or unsubstituted aryl, C₂-C₄₀ substituted or unsubstituted alkyleneoxy; C₃-C₄₀ substituted or unsubstituted alkyleneoxyalkyl; C₆-C₄₀ substituted or

unsubstituted alkylenearyl; C₆-C₃₂ substituted or unsubstituted aryloxy; C₆-C₄₀ substituted or unsubstituted alkyleneoxyaryl; C₆-C₄₀ oxyalkylenearyl, and mixtures thereof; or a cyclic orthoester having the formula:




wherein at least one R⁴ or R⁵ is derived from a fragrance raw material alcohol, each X is -C(R⁶)₂- wherein each R⁶ is independently hydrogen, C₁-C₂₂ linear or branched alkyl, C₂-C₂₂ linear or branched alkenyl, C₆-C₂₂ substituted or unsubstituted aryl, and mixtures thereof, Y is -CR⁷R⁸-, C=O, and mixtures thereof, wherein R⁷ and R⁸ are independently hydrogen, hydroxyl, nitro, nitrilo, C₁-C₃₀ substituted or unsubstituted linear alkyl, C₃-C₃₀ substituted or unsubstituted branched alkyl, C₃-C₃₀ substituted or unsubstituted cyclic alkyl, C₂-C₃₀ substituted or unsubstituted linear alkenyl, C₃-C₃₀ substituted or unsubstituted branched alkenyl, C₃-C₃₀ substituted or unsubstituted cyclic alkenyl, C₂-C₃₀ substituted or unsubstituted linear alkynyl, C₃-C₃₀ substituted or unsubstituted branched alkynyl, C₆-C₃₀ substituted or unsubstituted alkylenearyl, C₆-C₃₀ substituted or unsubstituted aryl, C₂-C₂₀ substituted or unsubstituted alkyleneoxy, C₃-C₂₀ substituted or unsubstituted alkyleneoxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₆-C₂₀ substituted or unsubstituted alkyleneoxyaryl, and mixtures thereof, or R⁷ and R⁸ can be taken together to form a spiroannulated ring or taken together with any R⁶ to form a fused ring, said spiroannulated or fused ring having from 3 to 8 carbons and optionally one or more heteroatoms in said ring, said ring further optionally substituted by one or more C₁-C₂₂ alkyl, C₁-C₂₂ alkenyl, C₆-C₁₂ aryl, C₆-C₂₂ alkylenearyl units, and mixtures thereof; m is from 0 to 14, p is from 0 to 14, and n is from 0 to 3; provided m + n + p is at least 1 and less than or equal to 14; and

B) optionally from about 1% by weight, a fragrance raw material component comprising:

- i) optionally at least 1% by weight, of a mixture of one or more base note fragrances;
- ii) optionally at least 1% by weight, of a mixture of one or more top or middle note fragrances;
- iii) optionally the balance carriers, fixatives, and other adjunct ingredients.

4. A composition according to any of Claims 1-3 wherein said aldehyde or ketone releasing pro-fragrance component releases a fragrance raw material selected from the group consisting of 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde, phenylacetaldehyde, methylnonyl acetaldehyde, 2-phenylpropan-1-al, 3-phenylprop-2-en-1-al, 3-phenyl-2-pentylprop-2-en-1-al, 3-phenyl-2-hexylprop-2-enal, 3-(4-isopropylphenyl)-2-methylpropan-1-al, 3-(4-ethylphenyl)-2,2-dimethylpropan-1-al, 3-(4-*tert*-butylphenyl)-2-methyl-propanal, 3-(3,4-methylenedioxyphenyl)-2-methylpropan-1-al, 3-(4-ethylphenyl)-2,2-dimethylpropanal, 3-(3-isopropylphenyl)butan-1-al, 2,6-dimethylhept-5-en-1-al, n-decanal, n-undecanal, n-dodecanal, 3,7-dimethyl-2,6-octadien-1-al, 4-methoxybenzaldehyde, 3-methoxy-4-hydroxybenz-aldehyde, 3-ethoxy-4-hydroxybenzaldehyde, 3,4-methylenedioxy-benzaldehyde, 3,4-dimethoxybenzaldehyde, and mixtures thereof.


5. A composition according to any of Claims 1-4 wherein said aldehyde or ketone releasing pro-fragrance component has the formula:




wherein R^8 is hydrogen, C_1 - C_{15} substituted linear alkyl, C_1 - C_{15} unsubstituted linear alkyl, C_1 - C_{15} substituted branched alkyl, C_1 - C_{15} unsubstituted branched alkyl, C_2 - C_{22} substituted or unsubstituted linear alkenyl, C_3 - C_{22} substituted or unsubstituted branched alkenyl, or mixtures thereof.

6. A composition according to any of Claims 1-5 wherein said aldehyde or ketone releasing pro-fragrance component is selected from the group consisting of:


a)



b)



c)




d) and mixtures thereof.


A composition according to any of Claims 1-6 further comprising at least 0.01% by weight, of a pro-accord which releases $n + 1$ fragrance raw materials wherein n is the number of fragrance raw materials from which said pro-accord is formed, n is from 1 to 3.

A composition according to any of Claims 1-7 wherein said β -amino pro-fragrance component comprises a G^2 unit which is methyl and a G^1 unit which is $-C(O)Y^1$ wherein Y^1 is selected from the group consisting of:


i) 2,6,6-trimethylcyclohex-2-enyl having the formula:




ii) 2,6,6-trimethylcyclohex-1-enyl having the formula:



- iii) 2-methylene-6,6-dimethylcyclohexanyl having the formula:



- iv) 2,6,6-trimethylcyclohex-3-enyl having the formula:



- v) and mixtures thereof.

10 9. A fragrance delivery system comprising:

- a) from 0.1% by weight, of a β -amino ketone pro-fragrance selected from the group consisting of 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(2-hydroxyethyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N-(5-hydroxy-3-oxapentyl)-1-butanone, and mixtures thereof;
- b) from 0.2% by weight, of an aldehyde releasing pro-fragrances selected from the group consisting of 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, and mixtures thereof;
- c) optionally from 0.1% by weight, of one or more alcohol releasing pro-fragrances or pro-accords;
- d) from 0.1% by weight, of one or more fragrance raw materials;
- e) optionally, from 1% by weight, of pre-blended perfume ingredients of fragrance raw material accords; and
- f) the balance carriers.

10. A composition according to Claim 9 wherein said alcohol releasing pro-fragrances or pro-accords are selected from the group consisting of tris-geranyl orthoformate, tris(*cis*-3-hexen-1-yl) orthoformate, tris(phenylethyl) orthoformate, tris(undecavertyl) orthoformate, bis(citronellyl) ethyl orthoacetate, tris(citronellyl) orthoformate, tris(*cis*-6-nonenyl) orthoformate, tris(phenoxyethyl) orthoformate, tris(geranyl, neryl) orthoformate (70:30 geranyl:neryl), tris(9-decenyl) orthoformate, tris(3-methyl-5-phenylpentanyl) orthoformate, tris(6-methylheptan-2-yl) orthoformate, tris([4-(2,2,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-yl] orthoformate, tris[3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-yl] orthoformate, trismenthyl orthoformate, tris(4-isopropylcyclohexylethyl-2-yl) orthoformate, tris-(6,8-dimethylnonan-2-yl) orthoformate, tris-phenylethyl orthoacetate, tris(*cis*-3-hexen-1-yl) orthoacetate, tris(*cis*-6-nonenyl) orthoacetate, tris-citronellyl orthoacetate, bis(geranyl) benzyl orthoacetate, tris(geranyl) orthoacetate, tris(4-isopropylcyclohexylmethyl) orthoacetate, tris(benzyl) orthoacetate, tris(2,6-dimethyl-5-heptenyl) orthoacetate, bis(*cis*-3-hexen-1-yl) amyl orthoacetate, and neryl citronellyl ethyl orthobutyrate, and mixtures thereof.

11. A composition according to either Claim 9 or 10 wherein said alcohol releasing pro-fragrances or pro-accords are selected from the group consisting of bis(ethyl) bis(geranyl) orthocarbonate, bis(ethyl) bis(phenylethyl) orthocarbonate, bis(ethyl) bis(*cis*-3-hexenyl) orthocarbonate, bis(ethyl) bis(citronellyl) orthocarbonate, bis(ethyl) bis(linalyl) orthocarbonate, bis(ethyl) bis(menthyl) orthocarbonate, bis(dodecyl) bis(geranyl) orthocarbonate, and bis(dodecyl) bis(phenylethyl) orthocarbonate, and mixtures thereof.

12. A fine fragrance or perfume comprising:
- from 0.1% by weight, of a β -amino pro-fragrance selected from the group consisting of 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(2-hydroxyethyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N-(5-hydroxy-3-oxapentyl)-1-butanone, and mixtures thereof;
 - from 0.2% by weight, of an aldehyde releasing pro-fragrances selected from the group consisting of 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-

00979492-11441


101
claim 9

oxazolidinecarboxylic acid methyl ester, 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)4-oxazolidinecarboxylic acid methyl ester, and mixtures thereof;


- c) optionally from 0.1% by weight, of one or more orthoester pro-accords; and
- d) the balance carriers.

5

13. A composition according to Claim 12 wherein said pro-accords have the formula:



wherein R is hydrogen, C₁-C₈ linear alkyl, C₄-C₂₀ branched alkyl, C₆-C₂₀ cyclic alkyl, C₆-C₂₀ branched cyclic alkyl, C₆-C₂₀ linear alkenyl, C₆-C₂₀ branched alkenyl, C₆-C₂₀ cyclic alkenyl, C₆-C₂₀ branched cyclic alkenyl, C₆-C₂₀ substituted or unsubstituted aryl, and mixtures thereof; R¹, R² and R³ are independently C₁-C₂₀ linear, branched, or substituted alkyl; C₂-C₂₀ linear, branched, or substituted alkenyl; C₅-C₂₀ substituted or unsubstituted cyclic alkyl; C₆-C₂₀ substituted or unsubstituted aryl, C₂-C₄₀ substituted or unsubstituted alkyleneoxy; C₃-C₄₀ substituted or unsubstituted alkyleneoxyalkyl; C₆-C₄₀ substituted or unsubstituted alkylenearyl; C₆-C₃₂ substituted or unsubstituted aryloxy; C₆-C₄₀ substituted or unsubstituted alkyleneoxyaryl; C₆-C₄₀ oxyalkylenearyl, and mixtures thereof; or a cyclic orthoester having the formula:



wherein at least one R⁴ or R⁵ is derived from a fragrance raw material alcohol, each X is -C(R⁶)₂- wherein each R⁶ is independently hydrogen, C₁-C₂₂ linear or branched alkyl, C₂-C₂₂ linear or branched alkenyl, C₆-C₂₂ substituted or unsubstituted aryl, and mixtures thereof, Y is -CR⁷R⁸-, C=O, and mixtures thereof, wherein R⁷ and R⁸ are independently hydrogen, hydroxyl, nitro, nitrilo, C₁-C₃₀ substituted or unsubstituted linear alkyl, C₃-C₃₀ substituted or unsubstituted branched alkyl, C₃-C₃₀ substituted or unsubstituted cyclic alkyl, C₂-C₃₀ substituted or unsubstituted linear alkenyl, C₃-C₃₀ substituted or unsubstituted branched alkenyl, C₃-C₃₀ substituted or unsubstituted cyclic alkenyl, C₂-C₃₀ substituted or unsubstituted linear alkynyl, C₃-C₃₀ substituted or unsubstituted branched alkynyl, C₆-C₃₀ substituted or unsubstituted alkylenearyl, C₆-C₃₀ substituted or

20

25

30

06979492-111401

101
claim 9

unsubstituted aryl, C₂-C₂₀ substituted or unsubstituted alkyleneoxy, C₃-C₂₀ substituted or unsubstituted alkyleneoxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₆-C₂₀ substituted or unsubstituted alkyleneoxyaryl, and mixtures thereof, or R⁷ and R⁸ can be taken together to form a spiroannulated ring or taken together with any R⁶ to form a fused ring, said spiroannulated or fused ring having from 3 to 8 carbons and optionally one or more heteroatoms in said ring, said ring further optionally substituted by one or more C₁-C₂₂ alkyl, C₁-C₂₂ alkenyl, C₆-C₁₂ aryl, C₆-C₂₂ alkylenearyl units, and mixtures thereof; m is from 0 to 14, p is from 0 to 14, and n is from 0 to 3; provided m + n + p is at least 1 and less than or equal to 14.

14. A composition according to either Claim 12 or 13 wherein said orthoester pro-accords release one or more fragrance raw material alcohols selected from the group consisting of 4-(1-methylethyl)cyclohexanemethanol, 2,4-dimethyl-3-cyclohexen-1-ylmethanol, (2,4-dimethylcyclohex-1-yl)methanol, (2,4,6-trimethyl-3-cyclohexen-1-yl)methanol, 2-phenylethanol, 1-(4-isopropylcyclohexyl)-ethanol, 2,2-dimethyl-3-(3-methylphenyl)propan-1-ol, 3-phenyl-2-propen-1-ol, 2-methyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-buten-1-ol, 3-methyl-5-phenylpentan-1-ol, 3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-ol, 2-methyl-4-phenylpentan-1-ol, *cis*-3-hexen-1-ol, 3,7-dimethyl-6-octen-1-ol, 3,7-dimethyl-2,6-octadien-1-ol, 7-methoxy-3,7-dimethyloctan-2-ol, 6,8-dimethylnonan-2-ol, *cis*-6-nonen-1-ol, 2,6-nonadien-1-ol, 4-methyl-3-decen-5-ol, benzyl alcohol, 2-methoxy-4-(1-propenyl)phenol, 2-methoxy-4-(2-propenyl)phenol, and mixtures thereof.
15. A composition according to any of Claims 1-14 wherein said orthoester pro-accord is a pro-accord which comprises *n* fragrance raw materials, said fragrance raw materials having a molecular weight greater than or equal to 100 g/mol and capable of releasing upon hydrolysis *n* + 1 fragrance raw materials, provided said pro-accord:
- has a molecular weight greater than or equal to 300 g/mol;
 - has a molecular weight at least two times greater than the lowest molecular weight fragrance raw material which comprises said pro-accord; and
 - has a fragrance release half-life of greater than or equal to 0.1 hours at pH 5.3 and less than or equal to 12 hours at pH 2.5 when measured in NaH₂PO₄ buffer; wherein *n* is an integer from 1 to 3.